



Moxifloxacin Use **Moxifloxacin** 3 times a day starting 3 days prior to surgery



Bromsite

Use **BromSite** 2 times a day starting 3 days prior to surgery.



Durezol

Do **NOT** use until after surgery

The SurgiCare Center of Idaho would like to make your visit to our surgical facility as pleasant as possible. We have provided you with a checklist of things which are important to do, or do not do the morning of your surgery.

- You will need to start both eye drops in your right/left eye on _____ before surgery. Instill one drop of each medication, waiting 5 minutes in between each drop. Do this as many times as directed above, up to day of surgery. **Surgery staff will administer drops on day of surgery.**

NOTHING TO EAT OR DRINK AFTER MIDNIGHT THE NIGHT BEFORE SURGERY

*****Please take blood pressure or seizure medications you are on with a small sip of water**.***

- You may wear your hearing aid or dentures.
-

- Please do not wear makeup the morning of the surgery. DO NOT wear any fragrances including cologne, perfume and lotion.
-

- Shower/Bathe before surgery, Wear clean, short sleeve, button-down front shirt. Bring dark sunglasses on day of surgery.
-

Arrive at _____ on _____ in surgery center.

1-Day Post Op at _____ on _____ in clinic.

You will not be able to drive immediately after surgery; therefore it will be necessary for you to make arrangements to have a family member or friend drive you home. Plan on being at our facility for about 3-4 hours.